

Cialis (Tadalafil)

It's available in four strengths: 2.5 mg, 5 mg, 10 mg, and 20 mg.

Dosage for benign prostatic hyperplasia (BPH) is 5mg daily.

1. Cialis should never be used by patients taking drugs containing Organic Nitrates such as: Minitran, Nitrek, Nitrodisc, Nitrogard, Nito-Dur, Nitrostat, Dilatrate, Nitroglyn, Nitrong, Nitrolingual spray, Deponit, Monoket, NitroPar, NitroTime, PentaerythritolTetranitrate, TransdermNitro, Ismo, Isordil, Sorbitrate, Imdur, Nitro-bid ointment, Erythatyl tetranitrate & Sodium Nitroprusside.

Substance such as Amyl nitrite (poppers), which are sometimes abused, should not be used with Cialis.

2. We recommend taking the drug 1 hour prior to intercourse.

3. Cialis will not cause an erection without sexual stimulation.

4. Common adverse reactions include: Headaches, flushing, diarrhea, upset stomach, dizziness, and rash.

5. Cialis should not be taken more than once a day.

Recommended dosage:

Take one pill an hour before intercourse, if it doesn't work after these attempts you may not be a candidate for this medication.

*The information provided on the site is for educational purposes only, and does not substitute for professional medical advice. Consult a medical professional or healthcare provider if you are seeking medical advice, diagnoses, or treatment.