

## **Levitra (Vardenafil)**

\*There have been reported deaths with the use of Levitra and there may be inherent dangers from the use of this drug. This is especially true in patients with heart disease & high blood pressure.

1. Levitra should never be used by patients taking drugs containing Organic Nitrates such as: Minitran, Nitrek, Nitrodisc, Nitrogard, Nito-Dur, Nitrostat, Dilatrate, Nitroglyn, Nitrong, Nitrolingual spray, Deponit, Monoket, Nitro-Par, Nitro-Time, Pentaerythritol Tetranitrate, Transderm-Nitro, Ismo, Isordil, Sorbitrate, Imdur, Nitro-bid ointment, Erytharyl tetranitrate, & Sodium Nitroprusside. Substance such as Amyl nitrite (poppers), which are sometimes abused, should not be used with Levitra.

2. We recommend taking the drug 1 hour prior to intercourse.

3. Levitra will not cause an erection without sexual stimulation.

4. Common adverse reactions include: Headaches, flushing, diarrhea, upset stomach, dizziness, and rash.

5. Levitra should not be taken more than once a day.

### **Recommended dosage:**

Take a 20 mg tablet one-hour prior to intercourse.

If it doesn't work after several attempts you may not be a candidate for this medication.

\*The information provided on the site is for educational purposes only, and does not substitute for professional medical advice. Consult a medical professional or healthcare provider if you are seeking medical advice, diagnoses, or treatment.