

Viagra (Sildenafil)

*There have been reported deaths with the use of Viagra and there may be inherent dangers from the use of this drug. This is especially true in patients with heart disease & high blood pressure.

1. Viagra should never be used by patients taking drugs containing Organic Nitrates such as: Minitran, Nitrek, Nitrodisc, Nitrogard, Nito-Dur, Nitrostat, Dilatrate, Nitroglyn, Nitrong, Nitrolingual spray, Deponit, Monoket, Nitro-Par, Nitro-Time, Pentaerythritol Tetranitrate, Transderm-Nitro, Ismo, Isordil, Sorbitrate, Imdur, Nitro-bid ointment, Erytharyl tetranitrate, & Sodium Nitroprusside.

Substance such as Amyl nitrite (poppers), which are sometimes abused, should not be used with Viagra.

2. Viagra will not cause an erection without sexual stimulation.

3. We recommend taking the drug 1 hour prior to intercourse.

4. Common adverse reactions include: Headaches, flushing, diarrhea, upset stomach, dizziness, rash & abnormal vision .

5. Viagra should not be taken more than once a day.

6. Avoid high fat meals (such as a cheeseburger & fries), it can make VIAGRA take longer to start working.

Recommended dosage:

Try a 50 mg for the first & second dose and if no erection increase to 2-50mg (100mg) pills for the remaining doses, if it doesn't work after these attempts you are not a candidate for this medication.

*The information provided on the site is for educational purposes only, and does not substitute for professional medical advice. Consult a medical professional or healthcare provider if you are seeking medical advice, diagnoses, or treatment.