




ANDRODERM

WHERE TO APPLY ANDRODERM[®]

There are 4 different areas where the Androderm[®] patch can be applied: the back, abdomen, upper arms, and thighs. Each day you're going to use a different spot than the previous day. Try not to reuse a site for at least 7 days. However, sometimes the area where you had your Androderm[®] patch may get red after the patch is removed. If the redness doesn't clear up, talk to your healthcare provider. He or she may suggest using an over-the-counter hydrocortisone cream.



Apply Androderm[®] patches to the areas in purple only.



SIMPLE TO APPLY

Before applying Androderm[®], make sure that the area you're applying it to is clean, dry, and has no broken skin.

1. Tear open (don't cut!) pouch and remove the patch, which has 2 tabs.
2. Peel the tabs apart to expose the adhesive and central reservoir area on the patch. Discard the protective liner and the silver disc.
3. Apply the patch to an area of skin that is clean, dry, and unbroken by placing it flat on the skin and firmly pressing it around the edges.



If the liner is hard to pull off, or if you see any glue or adhesive sticking to the liner, do not use the patch. Instead, throw it away and use a new one.

Make sure to avoid areas that are oily, perspire heavily, or are covered with hair, since Androderm[®] may not **stick well to these areas**. Do not apply Androderm[®] to your scrotum, buttocks, or over a bony area.

Androderm[®] is used to treat adult males who have low or no testosterone.

IMPORTANT SAFETY INFORMATION

You should not use Androderm[®] if you have breast cancer, known or suspected prostate cancer, or if you have pre-existing heart, kidney, or liver disease. You should also not use Androderm[®] if you are a pregnant or breastfeeding woman, as testosterone may cause fetal harm. If you have an enlarged prostate (BPH), you should be monitored for worsening signs and symptoms. Women and children should not be exposed to Androderm[®]. Taking testosterone may lead to a condition in which there are no sperm in the semen, and if you have risk factors, may cause sleep apnea (abnormal pauses in breathing). Your healthcare provider should monitor you for prostate specific antigen (PSA), liver function, lipids, and blood parameters. The most common adverse events in a study of 36 patients treated with Androderm[®] were skin irritation (itching at the application site), application site blisters, and back pain.

*The information provided on the site is for educational purposes only, and does not substitute for professional medical advice. Consult a medical professional or healthcare provider if you are seeking medical advice, diagnoses, or treatment.