UroPatient Podcast UroPatient.com Your go to place for Urologic Patient Education

Axiron (testosterone)

Axiron is a prescription medicine used to treat the symptoms of Testosterone Deficiency.

AXIRON can transfer from your body to others. This can happen if other people come into contact with the area where the AXIRON was applied. Signs of puberty that are not expected (for example, pubic hair) have happened in young children who were accidentally exposed to testosterone through skinto-skin contact with men using topical testosterone products like AXIRON. Women and children should avoid contact with the unwashed or unclothed area where AXIRON has been applied. If a woman or child makes contact with the application area, the contact area on the woman or child should be washed well with soap and water right away.

To lower the risk of transfer of AXIRON from your body to others, you should follow these important instructions:

Apply AXIRON only to your armpits

Wash your hands right away with soap and water after applying AXIRON
After the solution has dried, cover the application area with clothing. Keep the area covered until you have washed the application area well or have showered.

•If you expect another person to have direct skin-to-skin contact with your armpits, first wash the application area well with soap and water







PRIME THE PUMP

Always prime the pump when you open a new bottle of AURON. You only need to do this the first time you use it.

Gently press the pump 3 times into a sink or toilet. Do not use any of the liquid that came out when you primed.

Wash all the liquid three-uphy down the sink to avoid acuidential expressive to others.

IN THE BATHROOM

Apply AXIRDR at about the same time each moming after showering or bething. Make sure your underarms are clean and dry with no broken skin and apply it to the underarms enty.

Do sol apply AMRON to other pasts of your body like your alomach, penia, sorotum, shoelders, or upper arms.

PUMP THE RIGHT DOSE

Hold the nozzle over the applicator cup.

Genty press the pump once into the applicator cup. The applicator cup is made to hold 1 pump.

Avoid splashing AXINUM in the eyes. In case of contact with eyes, flush fhoroughly with water Himitation persists, seek medical advice.

WASH TO PREVENT TRANSFERENCE

Rinse the applicator cup with room-emperature running water, put I: dry with a lissue, and discard the tasse safely.

Turn the nocale so that it points away from the label, and place the applicator cop and cap back on the lexits. Store it safely cut of the reach of children. To receive child-resistant packaging, call 1-877-9-A2380M (4-877-421-438).

*The information provided on the site is for educational purposes only, and does not substitute for professional medical advice. Consult a medical professional or healthcare provider if you are seeking medical advice, diagnoses, or treatment.