

FLOMAX (TAMSULOSIN HCL)

IS INDICATED FOR THE TREATMENT OF THE SIGNS AND SYMPTOMS OF BENIGN PROSTATIC HYPERPLASIA (BPH). FLOMAX CAPSULES ARE NOT INDICATED FOR THE TREATMENT OF HYPERTENSION.

DOSAGE AND ADMINISTRATION

FLOMAX CAPSULES 0.4 MG ONCE DAILY IS RECOMMENDED AS THE DOSE FOR THE TREATMENT OF THE SIGNS AND SYMPTOMS OF BPH. IT SHOULD BE ADMINISTERED APPROXIMATELY **ONE-HALF HOUR FOLLOWING THE SAME MEAL EACH DAY** TO DECREASE THE RISK OF DIZZINESS.

FOR THOSE PATIENTS WHO FAIL TO RESPOND TO THE 0.4-MG DOSE AFTER TWO TO FOUR WEEKS OF DOSING, THE DOSE OF FLOMAX CAPSULES CAN BE INCREASED TO 0.8 MG ONCE DAILY. IF FLOMAX CAPSULES ADMINISTRATION IS DISCONTINUED OR INTERRUPTED FOR SEVERAL DAYS AT EITHER THE 0.4-MG OR 0.8-MG DOSE, THERAPY SHOULD BE STARTED AGAIN WITH THE 0.4-MG ONCE DAILY DOSE.

SIDE EFFECTS:

- LIGHT-HEADEDNESS OR DIZZINESS, ESPECIALLY WHEN CHANGING FROM SITTING TO STANDING HAS BEEN REPORTED.
- ABNORMAL EJACULATION ABNORMAL EJACULATION INCLUDES EJACULATION FAILURE, EJACULATION DISORDER, RETROGRADE EJACULATION AND EJACULATION DECREASE.
- ALLERGIC-TYPE REACTIONS SUCH AS SKIN RASH, PRURITUS SWELLING OF TONGUE, LIPS AND FACE AND ITCHING

YOU SHOULD INFORM YOUR OPHTHALMOLOGIST THAT YOU ARE TAKING THIS DRUG OR ANY OTHER PROSTATE DRUGS PRIOR TO EYE SURGERY.

*The information provided on the site is for educational purposes only, and does not substitute for professional medical advice. Consult a medical professional or healthcare provider if you are seeking medical advice, diagnoses, or treatment.